



Choosing the correct vaccine after a decision to offer a COVID-19 vaccine has been made

To support Ops, or HCPs making decisions about where to direct a patient, or vaccination centres who offer >1 vaccine each day



*** BE AWARE:** This is a tool to support clinicians choosing the correct vaccine, once the decision to vaccinate has been made. It does not consider vaccine eligibility (JCVI criteria, age group) or contraindications to all vaccinations (e.g. acutely unwell).

References

- Pg 17; Green Book Chapter 14a (01 July 2021)
- Pg 9; Green Book Chapter 14a (01 July 2021)
- Pg 16; Green Book Chapter 14a (01 July 2021)
- Pg 19; Green Book Chapter 14a (01 July 2021)
- History of thrombosis with thrombocytopenia syndrome is a 'Special warning and precaution for use' (AstraZeneca vaccine SPC; 19 July 2021)
- Pg 23; Green Book Chapter 14a (01 July 2021)
- NCL COVID Vaccine Pharmacy Group recommends avoiding AstraZeneca for patients with a history of thrombosis with thrombocytopenia syndrome whilst alternatives are available (15 July 2021)
- History of capillary leak syndrome is a 'Contraindication' (AstraZeneca vaccine SPC; 19 July 2021)
- Pg 21; Green Book Chapter 14a (01 July 2021)
- Pg 18; Green Book Chapter 14a (01 July 2021)
- Thrombosis with thrombocytopenia syndrome following vaccination with AstraZeneca is a 'Contraindication' (AstraZeneca vaccine SPC; 19 July 2021)
- London COVID-19 Clinical Advice Response Service (08 July 2021)
- Pg 17; Green Book Chapter 14a (30 July 2021)